



Home Infusion With Lumvoda

Home-based infusion care brings your intravenous (IV) medication treatment to your home and may offer scheduling considerations that some patients prefer.

A nurse from your selected home infusion provider will be available to support you throughout the process.



Preparing your medication

Setting up equipment

Administering your treatment

Supporting your treatment, which may include monitoring you during the infusion to help you feel supported during the infusion process

Please see Important Safety Information on [pages 7-8](#) and Full [Prescribing Information](#).



Determine if home infusion is right for you

Lumvoda™
veligrotug-vvze



If you are interested in home infusion, discuss the option with your doctor.



Once you and your doctor determine that home infusion is right for you, the next step is to check your insurance coverage. Enrollment in ViridianCares can help.



You don't need to be new to Lumvoda to qualify for home infusion

If you've already started treatment at an infusion center, hospital, or medical clinic, you still may be able to switch to home-based infusions for your remaining doses. Your doctor will determine whether home infusion is an appropriate option based on your medical needs and insurance coverage. In some cases, your doctor may determine that switching is appropriate.

If you're interested in home infusion, contact your doctor to inquire about whether it is right for you.

2

Prepare for your Lumvoa home infusion

Lumvoa™
veligrotug-vvze

After your appointment is scheduled, the home infusion company* will call you with details about your upcoming appointment. Use this call to ask any questions and make sure you know what to expect.

For the infusion itself, set up a comfortable chair where you can sit during your treatment, along with a clean, flat surface nearby where the nurse can prepare your medication and organize equipment.

Important questions to consider asking



Will Lumvoa be shipped to me before my appointment, or will the nurse bring it?

If shipped in advance, will I receive multiple doses at once or separate deliveries for each infusion?



Will the nurse bring all necessary infusion equipment, or should I expect supplies to arrive by mail?

Will infusion equipment stay in my home for the duration of my treatment, or will the nurse take it after each appointment?



Can I have a rolling stand to hold my infusion bag so I can move around or use the bathroom during the infusion?



What should I do to prepare my home for the infusion appointment?

*Home infusion services are provided by independent third-party providers selected by you and your healthcare team and are not associated with Viridian Therapeutics, Inc.

Please see Important Safety Information on [pages 7-8](#) and Full [Prescribing Information](#).

2

Prepare for your Lumvoa home infusion (continued)

Lumvoa™
veligrotug-vvze

Storing Lumvoa before your home infusion appointment

If your medication arrives at your home before your scheduled infusion, proper storage is essential.



Inside the shipping package, you'll find 1 or more cartons labeled "Lumvoa"



Keep the vial(s) inside the Lumvoa carton



Store the entire carton in your refrigerator at 36 °F to 46 °F (2 °C to 8 °C)



Keep it away from light



Do not freeze or shake



If you notice any damage to the Lumvoa carton when it arrives, contact your home infusion company right away.

Please see Important Safety Information on [pages 7-8](#) and Full [Prescribing Information](#).

3 What to expect with your infusion

Your treatment with Lumvoda will consist of 5 total infusions: 1 infusion every 3 weeks for 12 weeks.

Week 1	Weeks 3, 6, 9, & 12
 ~45 minutes	 ~30 minutes (if well tolerated) Your doctor may adjust how your treatment is administered based on how you tolerate each infusion.

On the day of an infusion, a nurse will arrive at your home, set up the infusion equipment, and prepare the infusion bag. Once the equipment and medication are ready, and you are seated in your chair, your nurse will insert a needle into your arm and start your Lumvoda infusion.

Your nurse may stay with you throughout the entire infusion. You're free to use the restroom or do relaxing activities in your chair like using a tablet, reading, or listening to music.

After each infusion is complete, your nurse may remain with you for a brief period to monitor for any reactions to Lumvoda.



Before the nurse leaves, confirm the date and time for your next infusion.

Please see Important Safety Information on [pages 7-8](#) and Full [Prescribing Information](#).



Questions?

VIRIDIAN[™]
cares

For more information,
or to speak with a ViridianCares
Patient Access Liaison (PAL), please call

1-866-VCARES1

Monday through Friday, 8 am to 8 pm ET



Please ensure your doctor has
submitted your Patient Enrollment
Form before reaching out for help.

Please visit [Lumvoa.com](https://lumvoa.com) for more resources that
may be helpful to you throughout your treatment journey.

Please see Important Safety Information on [pages 7-8](#)
and Full [Prescribing Information](#).

USE and IMPORTANT SAFETY INFORMATION



Use: Lumvoa is a prescription medicine used to treat Thyroid Eye Disease (TED), no matter if you've had TED for months or years.

What is the most important information I should know about Lumvoa?

Infusion reactions can happen during or soon after your infusion of Lumvoa. Tell your doctor or nurse right away if you have any of these symptoms during or after your treatment with Lumvoa:

- High blood pressure
- Headache
- Feeling hot
- Tiredness
- Chills

Lumvoa may worsen inflammatory bowel disease (IBD), such as Crohn's disease or ulcerative colitis. Tell your doctor right away if you have worsening IBD symptoms, which may include diarrhea with stomach pain or cramps, blood in your stools, sudden urgency to have a bowel movement, or feeling like you need to have a bowel movement even when there's little or no stool to pass.

Lumvoa may cause an increase in your blood sugar. Before starting treatment with Lumvoa, tell your doctor if you have high blood sugar or if you have diabetes. It is important for you to take your treatments and follow an appropriate diet for glucose control as prescribed by your doctor.

Lumvoa may cause severe hearing problems including hearing loss, which in some cases may be permanent. Tell your doctor if you have any signs or symptoms of hearing problems or changes in hearing.

USE and IMPORTANT SAFETY INFORMATION (continued)

Before receiving Lumvoa, tell your doctor if you:

- Have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- Have diabetes or know your blood sugar is high.
- Are pregnant or plan to become pregnant. Lumvoa may harm your unborn baby. Tell your doctor if you become pregnant or suspect you are pregnant during treatment with Lumvoa. Women who are able to become pregnant should use an effective form of birth control (contraception) prior to starting treatment, during treatment, and for 6 months after the final dose of Lumvoa.
- Are breastfeeding or plan to breastfeed. It is not known if Lumvoa passes into your breast milk. Talk to your doctor about the best ways to feed your baby during treatment with Lumvoa.
- All the medicines you take, including prescription and over-the-counter medicines, vitamins, and dietary and herbal supplements. Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

What are the possible side effects of Lumvoa?

The most common side effects of Lumvoa include muscle cramps or spasms, headache, hearing problems, high blood sugar, tiredness, diarrhea, ear discomfort, infusion reactions, nausea, cold-like symptoms, increases in a blood enzyme called creatine phosphokinase, dry skin, and high blood pressure.

This is not a complete list of all possible side effects. Tell your doctor or treatment team about any side effect you may have. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/safety/medwatch, or call 1-800-FDA-1088.

Lumvoa (500 mg) is an injection for infusion.

Please see Full [Prescribing Information](#).

© 2026 Viridian Therapeutics, Inc. All Rights Reserved. Lumvoa™ (veligrotug-vvze), Viridian Therapeutics®, ViridianCares™, and related trademarks are owned by Viridian Therapeutics, Inc. US-VG-0121 06/26

Lumvoa™
veligrotug-vvze

VIRIDIAN™
cares